



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

II. Solution by DR. O. E. GLENN, Drury College.

The theorem covering the general problem is due to Gauss,* and is the following: If p is any prime less than or equal to m , then the highest power of p dividing $m!$ is

$$p \left[\frac{m}{p} \right] + \left[\frac{m}{p^2} \right] + \left[\frac{m}{p^3} \right] + \dots = p \sum_{i=1}^{\infty} \left[\frac{m}{p^i} \right]$$

where $[s/t]$ stands for the greatest integer in s/t . Applying this we have $2^{500+250+125+62+31+15+7+3+1} = 2^{994}$, and similarly for the others.

Also solved by A. H. Holmes, and G. B. M. Zerr.

GEOMETRY.

263. Proposed by FREDERICK R. HONEY, Trinity College, Hartford, Conn.

Construct a sphere whose surface shall intersect the surface of any four given spheres in great circles.

Solution by G. W. GREENWOOD, M. A., McKendree College, Lebanon, Ill.

Let P be the center of a circle intersecting two circles, centers C, C' , in the extremities of diameters $AB, A'B'$, respectively. Draw through P a perpendicular to CC' , intersecting it in D . Then, if r be the radius of the intersecting circle, we have

$$r^2 = PC^2 + CA^2 = PC'^2 + C'A'^2.$$

$$\therefore PD^2 + DC^2 + CA^2 = PD^2 + DC'^2 + C'A'^2, \text{ and } DC^2 - DC'^2 = C'A'^2 - CA^2.$$

Hence D is a fixed point, and the locus of P is consequently a fixed line. By rotating the figure about CC' we find that the locus of the center of a sphere intersecting two given spheres in great circles is a certain plane.

Constructing these planes for three pairs of the given spheres, each sphere being involved, we get a common point as the center of the required sphere, assuming that the centers of the given spheres are not coplanar.

264. Proposed by B. F. FINKEL, A. M., Drury College, Springfield, Mo.

Let l and m be two straight lines intersecting in A . With A as center and any radius r describe a circle intersecting l and m in E, M and G, Q , respectively; and the bisector of the opposite angles formed by l and m in F and K . With I , the middle point of EA , as center, and radius, r , describe an arc intersecting the bisector of the opposite angles formed by l and m in O . With O as center, and radius $OA + r$ describe circle $FHCDBJF$; F and D the points of intersection of this circle with the bisector of opposite angle; H, B the intersections on l , and J, C on m . What is the ratio of arc HFJ to arc BD ?

**Disquisitiones Arithmeticae*.

Solution by G. B. M. ZERR, A. M., Ph. D., Parsons, W. Va.

Let 2θ be the angle formed by l, m . Also let A be the origin. Then $(\frac{1}{2}r\cos\theta, \frac{1}{2}r\sin\theta)$ are the coördinates of I where FD is the axis of abscissas.

$\therefore x^2 + y^2 - rx\cos\theta - ry\sin\theta = \frac{3}{4}r^2$ is the equation to the circle, center I .

$$\therefore y^2 - ry\sin\theta = \frac{3}{4}r^2 \text{ or}$$

$$y = \frac{1}{2}r\sin\theta \pm \frac{1}{2}\sqrt{(3r^2 + r^2\sin^2\theta)}.$$

$$\therefore OD = OA + r = y + r = r[1 + \frac{1}{2}\sin\theta + \frac{1}{2}\sqrt{(3 + \sin^2\theta)}].$$

Then if $\angle BOD = \phi$,

$$AO:BO = \sin(\phi - \theta):\sin\theta.$$

$$\therefore \sin(\phi - \theta) = \frac{[\sin\theta + \sqrt{(3 + \sin^2\theta)}] \sin\theta}{2 + \sin\theta + \sqrt{(3 + \sin^2\theta)}}$$

This gives ϕ . Now $\text{arc} HFJ:\text{arc} BD = 2\theta r:\phi DO$.

$$\therefore \text{arc} HFJ:\text{arc} BD = 2\theta:[2 + \sin\theta + \sqrt{(3 + \sin^2\theta)}]\phi.$$

265. Proposed by R. D. CARMICHAEL, Hartselle, Ala.

Find the Cartesian equation of a curve in a vertical plane such that a particle, sliding down the curve under the force of gravity alone, will require to pass from any point of beginning to the lowest point of the curve, a time proportional to the square of the distance to be traversed along the curve.

Solution by PROFESSOR WILLIAM HOOVER, Ph. D., Athens, Ohio.

With the usual notation, the equation of motion is

$$\frac{ds^2}{dt^2} = 2gy \dots\dots (1).$$

Let $t = ks^2$, or $s\sqrt{k} = \sqrt{t} \dots\dots (2)$. This gives

$$\frac{d^2s}{dt^2} = \frac{1}{4kt} = \frac{1}{4k^2s^2} = 2gy \dots\dots (3), \text{ or, } s = \frac{1}{2k\sqrt{2g}\sqrt{y}} \dots\dots (4).$$

$$\text{Differentiating, } ds = -\frac{1}{4k\sqrt{2g}} \cdot \frac{1}{y^{\frac{3}{2}}} \dots\dots (5). \quad \text{But } dx^2 + dy^2 = ds^2 \dots\dots (6).$$

Substituting (5) in (6), and arranging,

$$dx = \frac{\sqrt{(1 - 16k^2gy^3)}}{4k\sqrt{g}y^{\frac{3}{2}}} dy,$$

the differential equation of the curve, cartesian coördinates.

Also solved by G. B. M. Zerr. Professor Greenwood derives the *intrinsic* equation in the simple form $k^2 = 4s^2g\cos\phi$.

